

NATIONAL / INTERNATIONAL PRESENTATIONS

Infection rate in adult neurosurgical unit: Retrospective analysis of 501 patients. Associated factors, management, outcome and proposed modes of prevention.

L Hajipour, A D Mendelow, R P Sengupta, C J Gerber, A Jenkins, J J Nissen, N V Todd, P J Crawford, C Stokoe, Helen M Fernandes

Presented at Society of British Neurology Surgeons (SBNS), September 2002 (Middlesborough)

Hand washing rituals in theatre: Clean or dirty?

L Hajipour, L Longstaff, V Cleeve, N Brewster, P Henman

Presented at the;

- Scottish Orthopaedic annual meeting, Stirling (6th June 2003).
- British Trauma Society Annual Scientific Meeting, London (1st October 2003)

Poster presentation at the;

- Royal College of Surgeons of Edinburgh Clinical and Scientific Meeting (5th November 2003).
- 11th meeting of the combined Orthopaedic Associations, Sydney October 2004.

Significance of the tumour volume in pathological fracture.

L Hajipour, MR Acharya, WM Harper.

- Presented at the British Orthopaedic Oncology Association, June 2004.
- Poster presentation at EFORT, July 2005, Lisbon.

Metastatic Bone Disease: Are reamings useful?

L Hajipour, MR Acharya, WM Harper.

Presented at the;

- British Orthopaedic Oncology Association, June 2004.
- East Midlands Orthopaedic Research Meeting, November 2004.
- Naughton Dunn meeting, Birmingham November 2004

Poster presentation at the;

- EFORT, July 2005, Lisbon.

Attitudes of orthopaedic surgeons in the treatment of pathological fractures of long bones.

L Hajipour, MR Acharya, WM Harper.

Poster presentation at the;

- British Orthopaedic Oncology Association, June 2004.
- EFORT, July 2005, Lisbon.

Effect of tendon laceration and trimming on tendon friction coefficient along A2 pulley: In vitro study on turkey tendon.

Hajipour L, Dias J, Ullah A, Ahmed S, Aggarwal S, Gulihar A.

Presented at the:

Presentation at the;

- Naughton Dunn meeting, Stoke on Trent, May 2007.
- BSSH meeting, London , November 2007
- Presentation at the BOA, Liverpool, Sept 2008

E-poster in EFFORT, Nice, May 2008

Effect of early weight bearing on union rate following hindfoot arthrodesis: a retrospective study.

Hajipour L, Allen P E

Presentation at the;

- EFFORT, Nice, May 2008
- BOA, Liverpool, Sept 2008

Effect of different peripheral suture technique on the friction coefficient of the flexor tendon in Zone II. In vitro study on turkey tendons.

Hajipour L, Gulihar A, Dias J.

Presentation at the;

- East Midlands Nov 2008.
- Professor Harper Memorial Jan 2009.
- EFFORT 2009
- BOA Sept 2009

The Effect of Sequential Pulley Resection on the Coefficient of Friction when Intact and Following Flexor Tendon Repair in Zone 2: In vitro studies on paired turkey tendons.

L Hajipour, B Bhowal, N Taub, J Dias

Podium presentation at the BSSH, October 2012 (York).

Poster presentation IFSSH & IFSHT, New Delhi ,4th to 8th March 2013.

The effect of single suture knot on the coefficient of friction of Zone 2 intact flexor tendon: In vitro studies on paired turkey tendons.

L Hajipour, B Bhowal, N Taub, J Dias

Podium presentation in the forthcoming Triennial Congress of the IFSSH & IFSHT, New Delhi ,4th to 8th March 2013.

The effect of isolated running suture on the coefficient of friction of flexor tendon in Zone 2, in absence of tendon laceration: In vitro studies on turkey tendons.

L Hajipour, B Bhowal, N Taub, J Dias

Poster presentation IFSSH & IFSHT, New Delhi ,4th to 8th March 2013.

Contact surface Scanning Electron Microscopy (SEM)of flexor tendon and A2 pulley following tendon treatments. In vitro study on turkey tendons.

L Hajipour, B Bhowal, J Dias

Podium presentation IFSSH & IFSHT, New Delhi ,4th to 8th March 2013.

REGIONAL PRESENTATION

Total Hip Replacement in the UK; Does practice reflect guidelines?

Presented at the Northern Kreibich research session, Newcastle upon Tyne, November 2003.

Results: Key areas remain where there is a lack of consensus. These include consent, thromboprophylaxis, trainee supervision and discharge policy. Interestingly, the results suggest that around 30% of implants may not have a 10year NICE benchmark

Infection rate in adult Neurosurgical unit: Retrospective analysis of 501 patients. Associated factors, management, outcome and proposed modes of prevention.

Newcastle General Hospital, February 2002

Results: High rate of discitis, shunt and chest infection post-operatively, but results compare well with wound infection rate in the literature. Post-operative antibiotics show no benefit.

Analgesia for abdominal pain in Accident and Emergency Department; type, dose, route and timing of administration.

Sunderland Royal Hospital, Department of Accident and Emergency, August 2000.

Results: 70% of patients received appropriate analgesia within 2 hours of arrival to A/E.

Postoperative Transfusion Triggers – a prospective study, to evaluate transfusion criteria and up to date knowledge of transfusion medicine.

Ninewells Hospital, Dundee, PRHO Surgery, April, 1999.

Results: Poor knowledge of transfusion criteria among hospital doctors was noticed, leading to excessive use of blood transfusion services.

AUDIT PROJECT

Total Hip replacement in the UK – are guidelines reflected in practice?

Newcastle General Hospital, January 2003.

Results: Key areas remain where there is a lack of consensus. These include consent, thromboprophylaxis, trainee supervision and discharge policy. Interestingly, the results suggest that around 30% of implants may not have a 10year NICE benchmark

Infection rate in adult Neurosurgical unit: Retrospective analysis of 501 patients. Associated factors, management, outcome and proposed modes of prevention.

Newcastle General Hospital, February 2002.

Results: High rate of discitis, shunt and chest infection post-operatively, but results compare well with wound infection rate in the literature. Post-operative antibiotics show no benefit.

Analgesia for abdominal pain in Accident and Emergency Department; type, dose, route and timing of administration.

Sunderland Royal Hospital, Department of Accident and Emergency. August 2000

Results: 70% of patients received appropriate analgesia within 2 hours of arrival to A/E.

Postoperative Transfusion Triggers – a prospective study, to evaluate transfusion criteria and up to date knowledge of transfusion medicine.

Ninewells Hospital, Dundee, PRHO Surgery, April 1999.

Results: Poor knowledge of transfusion criteria among hospital doctors was noticed, leading to excessive use of blood transfusion services.

The effect of prophylactic antibiotics on the rate of pacemaker pocket infection.

Fourth year medical school Audit project, June 1996.

Results: High rate of Pacemaker pocket infection was detected in patients who did not received prophylactic antibiotics.

Effect of early weight bearing on union rate following hindfoot arthrodesis: a retrospective study.

Hajipour L, Allen P E

Early weight bearing following hindfoot arthrodesis has no adverse effect on the union rate.